

The homothetical-Hopf transformation of Pythagorean quadruples

Mircea Crasmareanu 

Faculty of Mathematics, “Al. I. Cuza”
700506, Iași, Romania
e-mail: mcrasm@uaic.ro

Received: 8 October 2025

Revised: 12 May 2026

Accepted: 17 May 2026

Online First: ... May 2026

Abstract: This note introduces a transformation of Pythagorean quadruples by using the composition between the Hopf map and a homothety of the space $\mathbb{R}^3$. Both the real algebra of complex numbers and the algebra of quaternions are used in this construction. Three examples are detailed, the first one concerning the well-known twin Pythagorean quadruple $(1, 2, 2, 3)$. The trigonometric parametrization of the Euclidean unit sphere $S^2 \subset \mathbb{E}^3$ allows us to prove that this transformation does not produce twin Pythagorean quadruples. A matrix approach for our transformation is also presented.

Keywords: Pythagorean quadruple, Hopf fibration, Homothety, Sphere.

2020 Mathematics Subject Classification: 11D09, 53C20, 11R52.

1 Pythagorean quadruples and their homothetical-Hopf transformation

For a natural number $n \geq 2$ let $\mathbb{E}^n := (\mathbb{R}^n, \langle \cdot, \cdot \rangle)$ be the n -dimensional *Euclidean space*:

$$\begin{cases} \langle \bar{x}, \bar{y} \rangle := x^1 y^1 + \dots + x^n y^n, & \|\bar{x}\|^2 := \langle \bar{x}, \bar{x} \rangle \geq 0, \\ \bar{x} = (x^1, \dots, x^n), & \bar{y} = (y^1, \dots, y^n). \end{cases} \quad (1.1)$$



For the real number $R > 0$ let $S^{n-1}(R)$ be the $(n - 1)$ -dimensional *sphere*:

$$S^{n-1}(R) := \{\bar{x} \in \mathbb{R}^n; \|\bar{x}\| = R\} = \{\bar{x} \in \mathbb{R}^n; \|\bar{x}\|^2 = R^2\} \quad (1.2)$$

with the usual *unit sphere* S^{n-1} when $R = 1$. Being a hypersurface in $\mathbb{E}^n$ the sphere $S^{n-1}(R)$ is itself a Riemannian manifold; in fact a *space form* since it has the constant sectional curvature $+1$.

The starting point of this short note is the well-known *Riemannian submersion*:

$$H : S^3 \subset \mathbb{C} \times \mathbb{C} \rightarrow S^2 \left(\frac{1}{2} \right) \subset \mathbb{C} \times \mathbb{R}, \quad H(z, w) := \left(z\bar{w}, \frac{1}{2}(|z|^2 - |w|^2) \right) \quad (1.3)$$

called *Hopf fibration*; see [6]. Here $\bar{w}$ denotes the complex conjugate of $w \in \mathbb{C}$.

Our interest consists of *Pythagorean quadruples*, namely $\square = (A, B, C, D) \in (\mathbb{N}^*)^4$ satisfying:

$$D^2 = A^2 + B^2 + C^2. \quad (1.4)$$

Denotes $\mathcal{PQ}$ their set; such a quadruple is called *primitive* if the greatest common divisor of its elements is 1. If $\mathbb{R}^4$ is endowed with the Minkowski-Lorentz product:

$$\langle \bar{x}, \bar{y} \rangle_{ML} := x^1 y^1 + \dots + x^3 y^3 - x^4 y^4$$

then $\square$ is a *null vector* with integer components. Let us call this fact as being *the physical interpretation* of the Pythagorean quadruples.

The relationship between Pythagorean quadruples and the Hopf map is provided by the quaternion algebra $\mathbb{H} := \mathbb{R}^4$ which is exactly the ambient space of S^3 with a special product; see again [6]. More precisely, the parametrization of any primitive $\square \in \mathcal{PQ}$ is ([7]):

$$\begin{cases} A := m^2 + n^2 - p^2 - q^2, B := 2(mq + np), C := 2(nq - mp), D := m^2 + n^2 + p^2 + q^2, \\ 0 < m, n, p, q \in \mathbb{N}, \quad 1 = \gcd(m, n, p, q), \quad m + n + p + q = \text{odd}. \end{cases} \quad (1.5)$$

Recall also that a non-zero real number λ defines the *homothety with a ratio λ* as:

$$h_\lambda^n : \mathbb{E}^n \rightarrow \mathbb{E}^n, \quad \bar{x} \rightarrow \lambda \bar{x}, \quad (1.6)$$

which implies that h_λ^n is a bijection map having the inverse $h_{1/\lambda}^n$. In fact, the set of all homotheties $\{h_\lambda^n\}$ acts on $\mathbb{E}^n$ as the commutative group $(\mathbb{R}^*, \cdot)$.

Usually, the study of Pythagorean quadruples is based on techniques from abstract algebra and number theory; now we return to the geometrical source of this notion through the above preparatory tools. Hence, the first main result of this work is:

Theorem 1.1. *Any Pythagorean quadruple $\square$ yields a new one:*

$$h_2^3 \circ H(\square) := (A^2, AC + BD, |AB - CD|, D^2) \in \mathcal{PQ}. \quad (1.7)$$

Proof. The use of $\mathbb{H}$ in producing the parametrization (1.5) inspires us to associate a quaternion $q(\square)$ as follows:

$$q(\square) := D + (A\bar{i} + B\bar{j} + C\bar{k}) = \text{Re}(q(\square)) + \text{Im}(q(\square)) \in \mathbb{H}. \quad (1.8)$$

The motivation of this choice is given by the fact that the square of $q(\square)$ in $\mathbb{H}$ is a *pure quaternion*:

$$\text{Re}((q(\square))^2) := D^2 - A^2 - B^2 - C^2 = 0 \quad (1.9)$$

from the definition of $\square$; a similar choice is performed in [4] when working with the plane $\pi : Ax + By + Cz + D = 0$ in $\mathbb{E}^3$. From the expression (1.8) we find two complex numbers which are useful for the application of the Hopf map:

$$\begin{cases} q(\square) = (D + A\bar{i}) + (B + C\bar{i})\bar{j}, \\ z(\square) := \frac{1}{D\sqrt{2}}(D + A\bar{i}) \in \mathbb{C}, \quad w(\square) := \frac{1}{D\sqrt{2}}(B + C\bar{i}) \in \mathbb{C}. \end{cases} \quad (1.10)$$

The fact that $\square$ is a Pythagorean quadruple assures that $(z(\square), w(\square))$ belongs to the sphere S^3 .

A direct computation of the composition $h_2^3 \circ H$ gives:

$$h_2^3 \circ H(z(\square), w(\square)) = \frac{1}{D^2}((AC + BD) + (AB - CD)\bar{i}, A^2) \in S^2, \quad (1.11)$$

which is the claimed conclusion. We point out that the direct algebraic computation confirms the validity of our use of the Hopf map:

$$\begin{aligned} (A^2) + (AC + BD)^2 + (AB - CD)^2 &= A^4 + A^2C^2 + B^2D^2 + A^2B^2 + C^2D^2 \\ &= A^2(A^2 + B^2 + C^2) + D^2(B^2 + D^2) \\ &= A^2D^2 + D^2(B^2 + C^2) \\ &= D^4. \end{aligned}$$

In order to explain also (1.5) we use the complex numbers $\alpha = \alpha(\square) := m + n\bar{i}$, $\beta = \beta(\square) := q + p\bar{i}$ to derive:

$$\square = (|\alpha|^2 - |\beta|^2, 2\text{Re}(\alpha\bar{\beta}), 2\text{Im}(\alpha\bar{\beta}), |\alpha|^2 + |\beta|^2). \quad (1.12)$$

This completes the proof. □

Definition 1.1. The map $H^2 : \mathcal{PQ} \rightarrow \mathcal{PQ}$ provided by the theorem will be called the *homothetical-Hopf transformation*.

Example 1.2 The *minimal* primitive Pythagorean quadruple is:

$$\begin{cases} \text{minimal} = (A = 1, B = C = 2, D = 3, m = n = q = 1, p = 0), \\ z(\text{minimal}) = \frac{1}{3\sqrt{2}}(3 + \bar{i}), \quad w(\text{minimal}) = \frac{\sqrt{2}}{3}(1 + \bar{i}) = \frac{2}{3}e^{\frac{\pi}{4}\bar{i}}, \\ \alpha(\text{minimal}) = \sqrt{2}e^{\frac{\pi}{4}\bar{i}}, \quad \beta(\text{minimal}) = 1 \in \mathbb{R}. \end{cases} \quad (1.13)$$

Sometimes a Pythagorean quadruple with $B = C$ is called a *twin*; see [8, p. 10]. We compute:

$$\begin{cases} h_2 \circ H(\text{minimal}) = \frac{1}{9}(8 - 4\bar{i}, 1), \quad H^2(\text{minimal}) = (1, 8, 4, 9), \\ \langle \text{minimal}, H^2(\text{minimal}) \rangle_{ML} = -2, \\ z(H^2(\text{minimal})) = \frac{1}{9\sqrt{2}}(9 + \bar{i}), \quad w(H^2(\text{minimal})) = \frac{2\sqrt{2}}{9}(2 + \bar{i}). \end{cases} \quad (1.14)$$

A new application of H^2 to $(1, 4, 8, 9) = (m = 1, n = q = 2, p = 0)$ gives the Pythagorean quadruple $(1, 44, 68, 81) = (m = 4, n = 5, p = 2, q = 6)$. Applying H^2 for the third time, we obtain $(1, 3632, 5464, 6561) = (m = 40, n = 41, p = 12, q = 56)$. $\square$

Remark 1.3 i) Denoting $\square = q(m, n, p, q)$ and $H^2(\square) = q(M, N, P, Q)$ it results:

$$\begin{cases} M = p^2 + q^2, & N = m^2 + n^2, & \alpha(H^2(\square)) = |\beta(\square)|^2 + |\alpha(\square)|^2 i, \\ (M^2 + N^2)Q = M(mq + np + mp - nq) + N(mq + np + nq - mp). \end{cases} \quad (1.15)$$

ii) An important map in the study of quaternions is *the conjugation*:

$$q \rightarrow \bar{q} := \text{Re}(q) - \text{Im}(q). \quad (1.16)$$

The effect of this map on the quaternion $q(\square)$ from (1.8) means the following transformation:

$$z(\square) \rightarrow \overline{z(\square)} \text{ (in } \mathbb{C}), \quad w(\square) \rightarrow -w(\square). \quad (1.17)$$

iii) Our Theorem 1.1. reveals a strong relationship between the Riemannian nature of the Hopf map and the pseudo-Riemannian nature of the Minkowski-Lorentz metric appeared in the physical interpretation of Pythagorean quadruples. It follows in this way the two faces of a coin titled “Pythagorean quadruple”. $\square$

Example 1.4 Let us consider now a non-twin Pythagorean quadruple. For example:

$$\square = (3, 6, 2, 7), \quad m = p = q = 1, n = 2 \quad (1.18)$$

gives: $H^2(\square) = (9, 48, 4, 49)$ corresponding to $M = Q = 2, P = 4$ and $N = 5$.

Example 1.5 A very interesting class of Pythagorean quadruples is produced by Pythagorean triples via the Exercise 3.17 from [9, p. 73]: If $\triangle := (u, v, w) \in (\mathbb{N}^*)^3$ is a *Pythagorean triple*, then:

$$(uv)^4 + (vw)^4 + (wu)^4 = (w^4 - u^2v^2)^2. \quad (1.19)$$

It follows that a proper $\triangle$ (for which we suppose $u < v < w$) yields the Pythagorean quadruple:

$$\square(\triangle) := (A = (wu)^2, B = (vw)^2, C = (uv)^2, D = w^4 - u^2v^2 = w^4 - A). \quad (1.20)$$

Concretely, the classical $\triangle_{\min} = (3, 4, 5)$ gives:

$$\begin{cases} \square(\triangle_{\min}) = (A = 15^2 = 225, B = 20^2 = 400, C = 12^2 = 144, D = 13 \cdot 37 = 481), \\ m = 8, \quad n = 17, \quad p = q = 16. \end{cases} \quad (1.21)$$

We leave it as an exercise to compute the image H^2 of this huge Pythagorean quadruple.

2 The special case of twin Pythagorean quadruples

First of all, we point out that if the point $P(x_0, y_0, z_0)$ of the unit sphere S^2 satisfies $y_0 = z_0$, then P also belongs to the ellipse:

$$E : x^2 + 2y^2 = 1 \quad (2.1)$$

with the eccentricity $e = \frac{1}{\sqrt{2}}$; this means that E is a *self-complementary ellipse*, a special class of ellipses studied in [3]. The plane $\pi : y - z = 0$ in the Euclidean space $\mathbb{E}^3$ is the plane through the origin $O(0, 0, 0)$ and having the unit normal vector $\bar{N} = e(0, 1, -1) \in S^2$.

Secondly, for a twin quadruple we have the complex number:

$$w(\square) = \frac{B}{D} e^{\frac{\pi}{4}i}. \quad (2.2)$$

Thirdly, the homotetical-Hopf transformation does not preserve the twin property (which reads as $m(q + p) = n(q - p)$) since for such a Pythagorean quadruple its image is:

$$H^2(\square) = (A^2, B(D + A) > B(D - A), D^2). \quad (2.3)$$

Fourthly, we use the well-known parametrization of the sphere S^2 with the angles $\varphi, \theta \in [0, \frac{\pi}{2})$ to express the first three components of the Pythagorean quadruple $\square$:

$$A = D \sin \varphi, \quad B = D \cos \varphi \cos \theta, \quad C = D \cos \varphi \sin \theta. \quad (2.4)$$

The twin property gives $\theta = \frac{\pi}{4}$. By denoting $\square = \square(\varphi, \theta)$ it follows H^2 as being a transformation on pairs of angles:

$$H^2 : (\varphi, \theta) \rightarrow (\Phi, \Theta). \quad (2.5)$$

A direct computation gives the relationships between these pairs of angles:

$$\begin{cases} \sin \Phi = \sin^2 \varphi \rightarrow 0 < \Phi < \varphi, \\ \tan \Theta = \frac{|\sin \varphi - \tan \theta|}{1 + \sin \varphi \tan \theta}. \end{cases} \quad (2.6)$$

The last relation provides a new proof of the third remark above; for $\theta = \frac{\pi}{4}$ it results:

$$\tan \Theta = \frac{1 - \sin \varphi}{1 + \sin \varphi} < 1. \quad (2.7)$$

The degenerate case $\varphi = 0$ (i.e., $A = 0$) allowing $\tan \Theta = 1$ yields the Diophantine equation $2B^2 = D^2$ with the degenerate solution $(A = B = C = 0 = D)$.

If we allow some of the components (A, B, C) to be negative, then we can find a twin image $H^2(\square)$ starting from a rational pair $(\sin \varphi, \tan \theta) \in \mathbb{Q} \times \mathbb{Q}$ only through the cases:

- 1) $\sin \varphi_1 = -\frac{1}{2}$, $\tan \theta_1 = \frac{1}{3}$; hence $\varphi_1 = -\frac{\pi}{6}$ and $\theta_1 = 18.43^\circ$;
- 2) $\sin \varphi_2 = -\frac{1}{3}$, $\tan \theta_2 = \frac{1}{2}$; hence $\varphi_2 = -19.47^\circ$ and $\theta_2 = 26.57^\circ$ giving $\theta_1 + \theta_2 = \frac{\pi}{4}$.

In the first case from $D = -2A$ it results $(2B)^2 + (2C)^2 = 3A^2$ with the only integer solution $(0, 0, 0)$ while the second case gives the Pythagorean quadruple $\lambda(-1, 2, 2, 3)$ which, following Example 1.2, does not produce a twin one. In conclusion, we prove the second main result of this study:

Theorem 2.1. *The image of the map H^2 on the subset of proper Pythagorean quadruples does not contain twin Pythagorean quadruples.*

3 The matrix approach to H^2

A second motivation for the choice of notation H^2 is that the homotetical-Hopf transformation can be thought as a *quadratic* one. There exist *linear* transformations on $\mathcal{PQ}$, see [8]. Recall that a similar fact holds for Pythagorean triples; see [1] or [2].

In this section, we treat the transformation H^2 with the tools of the matrix calculus. More precisely, supposing the order $0 < A < B < C < D$, we have

$$H^2(\square) = (A^2, AC + BD, CD - AB, D^2),$$

which means:

$$H^2(\square) = (A, B, C, D) \cdot M(\square), \quad M(\square) := \begin{pmatrix} A & \frac{C}{2} & -\frac{B}{2} & 0 \\ 0 & \frac{D}{2} & -\frac{A}{2} & 0 \\ 0 & \frac{A}{2} & \frac{D}{2} & 0 \\ 0 & \frac{B}{2} & \frac{C}{2} & D \end{pmatrix} \in M_4(\mathbb{R}). \quad (3.1)$$

The 4×4 matrix $M(\square)$ has the trace and the determinant as functions only of A and D :

$$\text{Tr}(M(\square)) = A + \frac{3D}{2}, \quad \det M(\square) = \frac{AD}{4}(A^2 + D^2) \quad (3.2)$$

and its characteristic polynomial is:

$$P_\lambda(\square) = (\lambda - A)(\lambda - D) \left[\lambda^2 - D\lambda + \frac{A^2 + D^2}{4} \right], \quad (3.3)$$

which implies that the eigenvalues of the $M(\square)$ are both real and complex:

$$0 < \lambda_1(\square) = A < \lambda_2(\square) = D \in \mathbb{R}, \quad \lambda_{3,4}(\square) = \frac{1}{2}(D \pm A\bar{i}) \in \mathbb{C}, \quad \sqrt{2}\lambda_3(\square) = D \cdot z(\square). \quad (3.4)$$

Also, the middle 2×2 block of the matrix $M(\square)$ suggests the special orthogonal matrix:

$$\frac{1}{\sqrt{A^2 + D^2}} \begin{pmatrix} D & -A \\ A & D \end{pmatrix} = \begin{pmatrix} \cos \psi & -\sin \psi \\ \sin \psi & \cos \psi \end{pmatrix} \in SO(2), \quad \psi = \psi(\square). \quad (3.5)$$

In fact, the occurrence of the matrix $M(\square)$ is natural if we consider the differential point of view. Namely, treating H^2 as a smooth map from $\mathbb{R}^4$ to $\mathbb{R}^4$, its Jacobian matrix of the partial derivatives is:

$$\begin{pmatrix} 2A & 0 & 0 & 0 \\ C & D & A & B \\ -B & -A & D & C \\ 0 & 0 & 0 & 2D \end{pmatrix},$$

which is the double of the transposed matrix $M(\square)^T$.

Example 3.1 Revisiting Example 1.2 we get:

$$\begin{cases} \text{Tr}(\text{minimal}) = \frac{11}{2}, & \det(\text{minimal}) = \frac{15}{2}, \\ \cos \psi(\text{minimal}) = \frac{3}{\sqrt{10}}, & \sin \psi(\text{minimal}) = \frac{1}{\sqrt{10}}, \end{cases} \quad (3.6)$$

and we point out that the angle $\frac{\pi}{2} - \psi(\text{minimal}) = 71.57^\circ$ is exactly the angle t_2 from Theorem 3.1 from [5].

It is worth to remark that the fourth degree polynomial $P_\lambda(\square)$ from (3.3) defines a plane quartic curve:

$$\mathcal{C}(\square) : y^2 = P_x(\square). \quad (3.7)$$

For our example we get:

$$\mathcal{C}(\text{minimal}) : y^2 = (x-1)(x-3) \left(x^2 - 3x + \frac{5}{2} \right) \quad (3.8)$$

which contains four integer points: $(1, 0), (3, 0), (5, \pm 10)$. $\square$

4 Conclusions

In this paper, the composition between the Hopf map and a homothety of the space $\mathbb{R}^3$ is used to produce new Pythagorean quadruples from old ones. The famous twin Pythagorean quadruple $(1, 2, 2, 3)$ serves as the first example. We prove that this transformation does not result in twin Pythagorean quadruples by using the trigonometrical parametrization of the unit sphere S^2 . In addition to Euclidean and Riemannian geometry, the elementary tools of linear algebra are considered in order to obtain a better image of this very interesting subject.

If we are looking for a possible generalization, we consider the general quadratic Diophantine equation:

$$(x^1)^2 + \dots + (x^n)^2 = (x^{n+1})^2.$$

It follows a rational point $\left(\frac{x^1}{x^{n+1}}, \dots, \frac{x^n}{x^{n+1}} \right) \in S^{n-1}$ and we point out that the Hopf map admits extensions to higher dimensions:

$$S^7 \rightarrow S^4, \quad S^{15} \rightarrow S^8.$$

The total spaces of these bundles, namely S^7 and S^{15} are the unit spheres in the Euclidean spaces $\mathbb{E}^8$ and $\mathbb{E}^{16}$, which means the occurrence of the algebra of octonions and sedenions, respectively, as successive members to $\mathbb{H}$ in the iterated Cayley–Dickson process. It remains an open problem what are their possible applications to the given Diophantine equation.

Acknowledgements

The author thanks the two anonymous referees for the many helpful comments which have improved the first version of the paper.

References

- [1] Crasmareanu, M. (2002). A new method to obtain Pythagorean triple preserving matrices. *Missouri Journal of Mathematical Sciences*, 14(3), 149–158.

- [2] Crasmareanu, M. (2019). Conics from symmetric Pythagorean triple preserving matrices. *International Electronic Journal of Geometry*, 12(1), 85–92.
- [3] Crasmareanu, M. (2021). Magic conics, their integer points and complementary ellipses. *Annals of the "Alexandru Ioan Cuza" University of Iași*, 67(1), 129–148.
- [4] Crasmareanu, M. (2025). A quaternionic product of planes in $\mathbb{E}^3$. *Romanian Journal of Mathematics and Computer Science*, 15(1), 9–14.
- [5] Crasmareanu, M., & Munteanu, M. (2025). The circular sum of points of an affine segment. *Turkish Journal of Mathematics*, 49(4), 411–420.
- [6] Cushman, R. H., & Bates L. M. (2015). *Global Aspects of Classical Integrable Systems* (2nd edition). Basel: Birkhäuser/Springer.
- [7] Frisch, S., & Vaserstein, L. (2012). Polynomial parametrization of Pythagorean quadruples, quintuples and sextuples. *Journal of Pure and Applied Algebra*, 216(1), 184–191.
- [8] Rushall, J., Gutierrez, M., & McCarty, V. (2020). On the complete tree of primitive Pythagorean quadruples. *Integers*, 20, Article ID A73.
- [9] Takloo-Bighas, R. (2018). *A Pythagorean Introduction to Number Theory. Right Triangles, Sums of Squares, and Arithmetic*. Undergraduate Texts in Mathematics, Cham: Springer.