

Para-mixed linear spaces

Mircea Crasmareanu

Faculty of Mathematics

University "Al. I. Cuza", 700506, Iași, România

<http://www.math.uaic.ro/~mcrasm>

email: mcrasm@uaic.ro

Abstract. We consider the paracomplex version of the notion of mixed linear spaces introduced by M. Jurchescu in [4] by replacing the complex unit i with the paracomplex unit j , $j^2 = 1$. The linear algebra of these spaces is studied with a special view towards their morphisms.

Introduction

It is well-known that up to isomorphisms there are three 2-dimensional real algebras: $\mathbb{C} = \mathbb{R}[X]/(x^2 + 1)$, $\mathbb{A} = \mathbb{R}[X]/(x^2 - 1)$, $\mathbb{D} = \mathbb{R}[X]/(x^2)$. The theory of the first algebra is richer than the other two, a fact corresponding to the field property of \mathbb{C} . Similar to the complex case, the paracomplex algebra \mathbb{A} has the basis $\{1, j\}$ with $j^2 = 1$; therefore the elements of \mathbb{A} are $z = x + jy$ with x and y real numbers. For historical details about the paracomplex algebra please see the survey [3].

Similar to the linear complex geometry there exists a paracomplex version as follows: let V be a real linear space. A *paracomplex structure* on V is an involution $J : V \rightarrow V$, $J^2 = 1_V$, such that the eigenspaces $V_{\pm} := \ker(1_V \pm J)$ have the same dimension. The pair (V, J) is then called a *paracomplex linear space*. If the hypothesis regarding the eigenspaces is dropped then we obtain the notion of *almost paracomplex structure*. An \mathbb{A} -linear map between the (almost) paracomplex linear spaces (V, J) and (V', J') is a linear map $T : V \rightarrow V'$ satisfying $T \circ J = J' \circ T$. Just like complex vector spaces are vector spaces over the field \mathbb{C} , the almost paracomplex linear spaces are free modules over \mathbb{A} .

2010 Mathematics Subject Classification: 15A03, 15A04

Key words and phrases: paracomplex structure, para-mixed linear space, morphism

In a series of papers ([4]-[6]) M. Jurchescu defined and uses the notions of *mixed linear space* and *mixed manifold* having as (local) model the direct product $\mathbb{R}^m \times \mathbb{C}^n$. The mixed manifolds as differentiable families of complex spaces are useful at the cross-road of complex analysis and complex geometry, for example regarding the smoothly parameterized Čech cohomology of complex manifolds, [1]. Here, following his ideas and restricted at the linear level we define the concept of *para-mixed linear space* by replacing \mathbb{C} with \mathbb{A} . A lot of properties of algebraic nature are similar to these frameworks and we hope to use further the notions considered now.

The paper is structured in two sections. The first one is devoted to the general theory of para-mixed linear spaces including the adapted linear maps between them and also their subspaces. A special attention is dedicated to the finite-dimensional case. The second section treats special morphisms between para-mixed linear spaces and the notion of *paracomplexification* of such spaces.

1 Para-mixed linear spaces and subspaces

Definition 1 *A (almost) para-mixed linear space is a triple (E, E_p, P) where E is a real linear space and E_p is a linear subspace of E endowed with an (almost) paracomplex structure P . Then E_p is called the paracomplex part of E while the quotient real linear space $E_r = E/E_p$ is the real part of E . The para-mixed space is pure real if $E_p = 0$ (i.e. $E_r = E$) and pure paracomplex if $E_r = 0$ (i.e. $E_p = E$). In the following we place always in the "almost" case and for simplicity we will drop this epithet.*

Example 1 *i) The fundamental example is $E = E_r = \mathbb{R}^m \times \mathbb{A}^n$ with $E_p = \mathbb{A}^n$ and $E_r = \mathbb{R}^m$. Hence, a para-mixed linear space can be thought as a (trivial) vector bundle over E_r with paracomplex fibres E_p ; also, para-mixed linear spaces can be thought as linear families of paracomplex spaces. Let us point out that vertical bundles endowed with paracomplex structures are recently studied in [2] and the geometry of polynomial sub-endomorphisms on a pair of distributions for a given manifold are studied in [7].*

ii) The paracomplex linear spaces will be considered as pure paracomplex para-mixed linear spaces while the real linear spaces will be considered as pure real para-mixed linear spaces. A para-mixed linear space E is simultaneous pure real and pure paracomplex if and only if $E = \{0\}$.

Definition 2 *A linear map $T \in L_{\mathbb{R}}(E, F) := L(E, F)$ between two para-mixed linear spaces is called a morphism if $T(E_p) \subseteq F_p$ and the induced map $T_p =$*

$T|_{E_p} : E_p \rightarrow F_p$ is an \mathbb{A} -linear map. T is called an antimorphism if $T(E_p) \subseteq F_p$ and T_p is an \mathbb{A} -antilinear map: $T(x + jy) = Tx - jTy$. T_p is the paracomplex part of T while $T_r = T|_{E_r} : E_r \rightarrow F_r$ is the real part of T . Denotes by $\text{Hom}(E, F)$ the set of all morphisms from E to F and $T \in \text{Hom}(E, F)$ is called isomorphism if both T_p and T_r are bijective maps.

Remark 1 *i) If E is a pure real para-mixed linear space and $T \in L(E, F)$ then T is both morphism and antimorphism. If F is a pure real para-mixed linear space and $T \in L(E, F)$ then T is a morphism if and only if $T_p = 0$. If E is a general para-mixed linear space then $(1_{E_r}, P) \in \text{Hom}(E, E)$ by considering the decomposition $E = E_r \otimes E_p$ (see also the Corollary 1 below).*

ii) The class of para-mixed linear spaces with their morphisms defines a category which contains the category of real linear spaces as well as the category of paracomplex linear spaces. The consideration of the paracomplex part (for spaces and morphisms) is a functor from the category of para-mixed linear spaces to the category of paracomplex linear spaces and similar for the consideration of the real part (for spaces and morphisms).

iii) Fix E and F two para-mixed linear spaces and $T_1, T_2 \in L(E, F)$ two (anti)morphisms. Consider also two real numbers α, β . It follows that $\alpha T_1 + \beta T_2$ is also an (anti)morphism with $(\alpha T_1 + \beta T_2)_i = (\alpha T_1)_i + (\beta T_2)_i$ for $i \in \{p, r\}$.

iv) The linear map $(x, y) \in \mathbb{R}^2 \rightarrow z = x + jy \in \mathbb{A}$ is a bijective morphism which is not an isomorphism.

v) $\text{Hom}(E, F)$ is a para-mixed linear space with the paracomplex part

$\text{Hom}(E, F)_p = \{T \in \text{Hom}(E, F); T_r = 0\}$ and real part $\text{Hom}(E, F)_r = \text{Hom}(E, F_r) \simeq \text{Hom}(E_r, F_r)$. The corresponding paracomplex structure is the map

$T \in \text{Hom}(E, F) \rightarrow T \circ (1_{E_r}, P) \in \text{Hom}(E, F)$.

vi) Fix $T \in \text{Hom}(E, F)$ and let H be another para-mixed linear space. Let $T_ : \text{Hom}(H, E) \rightarrow \text{Hom}(H, F)$ be the composition with T at left and $T^* : \text{Hom}(F, H) \rightarrow \text{Hom}(E, H)$ be the composition with T at right. Then T_* and T^* are morphisms with respect to the para-mixed structure from v).*

A first structural result is provided by:

Proposition 1 *In the category of para-mixed linear spaces a given para-mixed linear space E is isomorphic with the direct product $E_r \times E_p$.*

Proof. We have the canonical maps: $i : E_p \rightarrow E$ and $\pi : E \rightarrow E_r$. There exists the maps $q : E \rightarrow E_p$ and $\rho : E_r \rightarrow E$ such that: $q \circ i = 1_{E_p}$, $i \circ q + \rho \circ \pi = 1_E$. It follows that these maps i, π, q, ρ are morphisms of para-mixed linear spaces and $(\rho, i) : E_r \times E_p \rightarrow E$ is an isomorphism. ■

It follows directly:

Corollary 1 *Let $E = E_r \times E_p$ and $F = F_r \times F_p$ be para-mixed linear spaces and $T \in L(E, F)$. Then $T \in \text{Hom}(E, F)$ if and only if it has the expression:*

$$T = \begin{pmatrix} T_2 & 0 \\ \alpha & T_1 \end{pmatrix} \quad (1.1)$$

where $T_1 : E_p \rightarrow F_p$ is a \mathbb{A} -linear map while $T_2 : E_r \rightarrow F_r$ and $\alpha : E_r \rightarrow F_p$ are real linear maps. In this decomposition, T_1 is the paracomplex part of T and T_2 is the real part of T .

A characterization of isomorphisms is provided by:

Proposition 2 *Let E, F and T as above. Then the following statements are equivalent:*

- i) T is an isomorphism,
- ii) T, T_1 and T_2 are all bijective,
- iii) two of the maps T, T_1 and T_2 are bijective.

A special study can be performed in finite-dimension:

Definition 3 *Let E be a para-mixed linear space and $m, n \in \mathbb{N}$. We say that E is of (m, n) -type if E_r is a real linear space of dimension m and E_p is a paracomplex space of dimension n . A frame on E is a set of vectors $B = \{e_1, \dots, e_m, e_{m+1}, \dots, e_{m+n}\}$ with $\{e_{m+1}, \dots, e_{m+n}\}$ a basis in E_p and $\{[e_1], \dots, [e_m]\}$ a basis in E_r where $[e]$ is the class of $e \in E$ considered in E_r .*

A characterization of this notion is:

Proposition 3 *Fix E a para-mixed linear space of (m, n) -type and $B = \{e_1, \dots, e_{m+n}\} \subset E$ with $\{e_{m+1}, \dots, e_{m+n}\} \subset E_p$. Then B is a frame on E if and only if the map:*

$$T : (x^1, \dots, x^{m+n}) \in E_f = \mathbb{R}^m \times \mathbb{A}^n \rightarrow x = x^i e_i \in E \quad (1.2)$$

belongs to $\text{Hom}(E_f, E)$.

Definition 4 *Let E be para-mixed linear space and W a subspace of E . Then W is called para-mixed subspace of E if $W \cap E_p$ is a paracomplex subspace of E_p which means that $x \in W \cap E_p$ implies $jx \in E_p$.*

A para-mixed subspace W will be considered itself as a para-mixed linear space with $W_p = W \cap E$. Hence, the inclusion map $i : W \rightarrow E$ is a morphism with i_p and i_r injective maps.

Example 2 *i) For $T \in \text{Hom}(E, F)$ the kernel $\ker T = T^{-1}(0_F)$ is a para-mixed subspace of E .*

ii) The intersection of an arbitrary family of para-mixed subspaces is again a para-mixed subspace.

iii) Let E be a pure real para-mixed linear space and $W \subset E$ a (real) subspace. Then W is a para-mixed subspace. A similar property holds for the paracomplex case.

2 Monomorphisms and epimorphisms

Definition 5 *Let $T \in \text{Hom}(E, F)$.*

i) T is called monomorphism if there exists a para-mixed linear space G and $R \in \text{Hom}(G, F)$ such that the map $(T, R) : E \times G \rightarrow F$ is an isomorphism.

ii) T is called epimorphism if there exists a para-mixed linear space G and $R \in \text{Hom}(E, G)$ such that the map $(R, T) : E \rightarrow F \times G$ is an isomorphism.

A characterization of these types of morphisms is given by:

Proposition 4 *Let $T \in \text{Hom}(E, F)$ be given.*

I) The following statements are equivalent:

- a) T is a monomorphism,*
- b) T and T_r are injective maps,*
- c) T_p and T_r are injective maps,*
- d) T have an inverse morphism at left.*

II) Also, the following statements are equivalent:

- e) T is an epimorphism,*
- f) T and T_p are surjective maps,*
- g) T_p and T_r are surjective maps,*
- h) T have an inverse morphism at right.*

Proof. a) \Rightarrow b). From hypothesis the maps (T, R) and $(T, R)_r : E_r \times G_r \rightarrow F_r$ is bijective and then T, T_r are injective. b) \Rightarrow c). It is obvious.

c) \Rightarrow d). Consider the decomposition (1.1) of T . Since T_p is injective it follows the existence of R_1 an \mathbb{A} -linear map which is inverse at left. Similar, from T_r

being injective it result the existence of R_2 a \mathbb{R} -linear map which is inverse at left. The map $R : F \rightarrow E$ given by:

$$R = \begin{pmatrix} R_r & 0 \\ -R_p \alpha R_r & R_p \end{pmatrix}$$

is a morphism from F to E with $R \circ T = \text{Id}_E$.

d) \Rightarrow a). Let $R : F \rightarrow E$ be the inverse at left of T and consider $G = \ker R$ together with the inclusion $i : G \rightarrow F$. From $R \circ (1_F - T \circ R) = 0$ it results the existence of $w \in \text{Hom}(F, G)$ such that $T \circ R + i \circ w = 1_F$. Then $w \circ i = 1_G$ and $w \circ T = 0$. Let $A = (T, i) : E \times G \rightarrow F$ and $B = (R, w) : F \rightarrow E \times G$. With the equations above it follows that A and B are isomorphisms with $B = A^{-1}$.

The equivalences from II are analogous. \blacksquare

Corollary 2 *Fix $T \in \text{Hom}(E, F)$. Then T is a monomorphism if and only if $T(E)$ is a para-mixed subspace of F and the induced map $T' : E \rightarrow T(E)$ is an isomorphism. Also, T is an epimorphism if and only if the co-induced map $T'' : E/\ker T \rightarrow F$ is an isomorphism.*

Proof. Suppose that T is a monomorphism. Since T_r is injective it results that $T(E) \cap F_p = T_p(E_p)$ and so, $T(E)$ is a para-mixed subspace in F . It follows also that $T' \in \text{Hom}(E, T(E))$ and its paracomplex part T'_p is surjective. From $T = \text{injective}$ we get that T' and T'_p are bijective maps and then T' is an isomorphism. Similar arguments hold for the second part. \blacksquare

Let us remark that an injective $T \in \text{Hom}(E, F)$ is not a-priori a monomorphism and the example is provided by the inclusion $\mathbb{R} \rightarrow \mathbb{A}$. In order to include this class we consider:

Definition 6 $T \in \text{Hom}(E, F)$ is called cartesian monomorphism if it is injective and for every para-mixed linear space G and every map $\alpha : G \rightarrow E$ we have that $T \circ \alpha$ is a morphism if and only if α is a morphism.

This notion is useful for another concept:

Definition 7 A paracomplexification of the para-mixed linear space E is a pair (E^p, ρ) with E^p a paracomplex linear space and $\rho \in \text{Hom}(E, E^p)$ injective and satisfying $\rho(E) + j\rho(E) = E^p$ and $\rho(E) \cap j\rho(E) = \rho(E_p)$.

A characterization of this notion is given by:

Theorem 1 *i) Every para-mixed linear space E have a paracomplexification.
 ii) The morphism ρ is a cartesian monomorphism.
 iii) A pair (E^P, ρ) is a paracomplexification of E if and only if E^P is a paracomplex linear space and the map $\rho^* : \text{Hom}(E^P, F) \rightarrow \text{Hom}(E, F)$ given in Remark 1(vi) is bijective for any paracomplex linear space F .*

Proof. i) Let $E = E_r \times E_p$ the canonical decomposition of E and consider the space:

$$E^P = (E_r \otimes_{\mathbb{R}} \mathbb{A}) \times E_p.$$

It results that E^P is a paracomplex linear space. One define the map $\rho : E \rightarrow E^P$ by $\rho(x_2, x_1) = (x_2 \otimes 1, x_1)$ and a straightforward computation gives that (E^P, ρ) is a paracomplexification of E .

ii) Let G be a para-mixed linear space and $\alpha : G \rightarrow E$ such that $\rho \circ \alpha \in \text{Hom}(G, E^P)$. Fix $z \in G_p$; then:

$$\rho \circ \alpha(z) = j(\rho \circ \alpha)(jz) \in \rho(E) \cap j\rho(E) = \rho(E_p)$$

and the injectivity of ρ yields that $\alpha(z) \in E_p$. Also, $\rho \circ \alpha(jz) = j(\rho \circ \alpha(z)) = \rho(j\alpha(z))$ and again the injectivity of ρ gives $\alpha(jz) = j\alpha(z)$. These facts together with the \mathbb{R} -linearity means that $\alpha \in \text{Hom}(G, E)$.

iii) Fix (E^P, ρ) a paracomplexification of E , F a paracomplex linear space and $u \in \text{Hom}(E, F)$. Define then $v : E^P \rightarrow F$:

$$v(\rho(x) + j\rho(y)) := u(x) + ju(y)$$

for all $x, y \in E$. Since $\rho(x) + j\rho(y) = \rho(x') + j\rho(y')$ if and only if $y' - y \in E_p$ and $x - x' = j(y' - y)$ it results that v is well defined. It follows that v is a \mathbb{A} -linear map and $v \circ \rho = u$. We get also the uniqueness of v with these two properties. \blacksquare

Example 3 *i) Let E be a real linear space endowed with the paracomplex structure J . Consider then $E^P = (E, J) \oplus (E, -J)$ and the diagonal map $\rho : E \rightarrow E^P$. Then (E^P, ρ) is a paracomplexification of E . Indeed, every vector $(e_1, e_2) \in E^P$ has a decomposition $(e_1, e_2) = (x + Jy, x - Jy)$ with $x, y \in E$. More precisely, $x = \frac{1}{2}(e_1 + e_2)$ and $y = \frac{1}{2}(Je_1 - Je_2)$.*

Let now F be a paracomplex linear space and $u : E \rightarrow F$ a \mathbb{R} -linear map. The unique \mathbb{A} -linear map $v : E^P \rightarrow F$ satisfying $v \circ \rho = u$ is:

$$v(e_1, e_2) = \frac{1}{2}(u(e_1) - ju(Je_1)) + \frac{1}{2}(u(e_2) + ju(Je_2)).$$

ii) Let E, F be para-mixed linear spaces and fix $\rho : F \rightarrow \mathbb{F}^p$ a paracomplexification of F . Then $\rho_* : \text{Hom}(E, F) \rightarrow \text{Hom}(E, \mathbb{F}^p)$ given in Remark 1(vi) is a cartesian monomorphism. In general, $(\text{Hom}(E, \mathbb{F}^p), \rho_*)$ is not a paracomplexification of $\text{Hom}(E, F)$ since $\text{Hom}(E, \mathbb{F}^p)$ is a paracomplex linear space satisfying $\rho_*(\text{Hom}(E, \mathbb{F}^p)) \cap j\rho_*(\text{Hom}(E, \mathbb{F}^p))$ but generally it do not satisfies $\rho_*(\text{Hom}(E, \mathbb{F}^p)) + j\rho_*(\text{Hom}(E, \mathbb{F}^p)) = (\rho_*(\text{Hom}(E, \mathbb{F}^p)))^p$.

References

- [1] T. Bailey; M. Eastwood; S. Gindikin, Smoothly parameterized Čech cohomology of complex manifolds, *J. Geom. Anal.*, **15** (2005), no. 1, 9–23. MR2132263 (2006b:32013)
- [2] M. Crasmareanu; L.-I. Pişcoran, Weak para-CR structures on vertical bundles, *Adv. Appl. Clifford Algebr.*, **26** (2016), no. 4, 1127–1136. MR3564462
- [3] V. Cruceanu; P. Fortuny; P. M. Gadea, A survey on paracomplex geometry, *Rocky Mountain J. Math.*, **26** (1996), no. 1, 83–115. MR1386154 (97c:53112)
- [4] M. Jurchescu, *Variétés mixtes*, in Romanian-Finnish Seminar on Complex Analysis (Proc., Bucharest, 1976), 431–448, Lecture Notes in Math., 743, Springer, Berlin, 1979. MR0552907 (80m:32011)
- [5] M. Jurchescu, *Espaces mixtes*, in "Complex analysis"-Fifth Romanian-Finnish Seminar, Part 2 (Bucharest, 1981), 37–57, Lecture Notes in Math., 1014, Springer, Berlin, 1983. MR0738113 (85j:32048)
- [6] M. Jurchescu, Variétés mixtes et cohomologie, *Riv. Mat. Univ. Parma*, (4) **10** (1984), special vol. 10, 55–79. MR0777315 (86e:32036)
- [7] F. Özdemir; M. Crasmareanu, Geometrical objects associated to a substructure, *Turkish J. Math.*, **35** (2011), no. 4, 717–728. MR2896417

Received: February 16, 2017